

Endodontic Treatment in Artificial Deciduous Teeth by Manual and Mechanical Instrumentation: A Pilot Study

¹Fernanda Hecksher, ²Bruno Vidigal, ³Patricia Coelho, ⁴Diassanam Otoni, ⁵Christiano Alvarenga, ⁶Eduardo Nunes

ABSTRACT

Introduction: “*In vitro*” studies seek to simulate actual biological conditions in the laboratory and providing principles to be studied later, to facilitate the development of new techniques of root instrumentation in primary teeth and to ensure the integrity and function of the element.

Aim: This study aimed to demonstrate the technological advances in endodontics by endodontic treatment performed on artificial primary teeth using a rotary instrumentation system and reciprocation.

Materials and methods: Instrumentation of the root canal was performed via a manual, rotatory and Reciproc system.

Results: The rotatory systems can facilitate endodontic treatment in one session.

Conclusion: These procedures become increasingly easy and rapid with the help of technological advances in dentistry.

Keywords: Deciduous teeth, Endodontics, Instrumentation.

How to cite this article: Hecksher F, Vidigal B, Coelho P, Otoni D, Alvarenga C, Nunes E. Endodontic Treatment in Artificial Deciduous Teeth by Manual and Mechanical Instrumentation: A Pilot Study. *Int J Clin Pediatr Dent*, 2018;11(6):510-512

Source of support: Nil

Conflict of interest: None

INTRODUCTION

The dental pulp of deciduous teeth can become involved much earlier than permanent teeth in advance of carious lesions. In addition, exposure of the pulp can also occur much more often during cavity preparation due to the thickness of the enamel and dentin, which are very often thinner in deciduous teeth. It is also noteworthy that traumatic injuries, especially in anterior teeth, occur frequently, contributing to a public health problem.^{1,2}

^{1-3,6}Professor, ^{4,5}Specialist

^{1-3,6}Department of Dentistry, PUC-Minas, Minas Gerais, Brazil

^{1,4,5}Department of Dentistry, São Leopoldo Mandic, Minas Gerais, Brazil

³Department of Dentistry, Fainor, Minas Gerais, Brazil

²Department of Dentistry, Newton Paiva, Minas Gerais, Brazil

Corresponding Author: Bruno Vidigal, Professor, Department of Dentistry, PUC-Minas, Minas Gerais, Brazil, e-mail: bruno.vidigal@newtonpaiva.br

The main goal of dentistry is to maintain the integrity and function of the primary dentition to its physiological exfoliation. When the deciduous pulp is compromised, endodontic treatment should be performed to preserve the integrity and function of the tooth and its supporting tissues.³ The instrumentation of the root canal may be performed with manual or mechanical tools.⁴

The instruments that incorporate reciprocating movements were introduced in the market to model the dental channel using only a file. These file instruments have a different mechanism compared with other previously developed files. The system is designed to be used with a reciprocating.⁵

Endodontic treatment with rotary systems can contribute significantly to reduce the duration of clinical care for pediatric patients. However, it is essential to evaluate the dentin wear created by both hands and by rotary instruments to analyze the safety of this procedure in the different root thirds and the presence of resorption.⁶ This study aims to demonstrate the potential applications of technological advances in endodontics in pediatric dentistry.

METHODOLOGY

The sample comprised artificial deciduous teeth with coronary and root pulp (Denarte, São Paulo, SP, Brazil) divided into three groups: group one was instrumented with a manual technique (G1), group 2 used a roundabout technique (G2) and group 3 used a reciprocating instrumented technique (G3).

The manual instrumentation group used instrumentation techniques such as the Scheduling Crown-Down Manual, which were performed with Kerr type files 1st series 21 mm (VDW, Munich, Germany). The instrumentation system used RECIPROC file Reciproc 25/0.08 (VDW, Munich, Germany), while the instrumentation group's rotary system utilized Mtwo files of sizes 10/0.4, 15/0.5, 20/0.6, 25/0.6–21 mm, adapted to the VDW engine (VDW, Munich, Germany).

The procedure to gain access to the root canal was performed by an endodontist experienced in crown-opening with a round burn^o 1011, exploitation of channels C file–Pilot # 10 0.2, 21 mm, to the working length (CT) without resistance, and widening the entrance of the conduits with Gates Glidden drills using #2–4. It was then performed at timed instrumentation conduits.

We used a stopwatch to measure the instrumentation time in each root canal.

RESULTS

The comparisons between the manual instrumentation group and the rotary and reciprocating systems showed that all the three groups were able to perform the instrumentation of the conduits. The average time required for the manual technique was 4.4 minutes, while with the rotary and reciprocating systems required 3.4 minutes (Graph 1).

CLINICAL APPLICATIONS

If we consider these procedures implemented in professional clinical practice for 4 weeks, from the 2nd to the 6th week, and with two patients per day, the rotary and reciprocating systems will have saved approximately 24% of time spent relative to the manual procedure (Graphs 2 and 3).

Considering the practice over the course of one year, at a rate of two patients per day, the time saving is still

approximately 24%. Thus, this economical approach could result in 18 more patients being seen per month, as indicated by the formula below.

$$1.8 \text{ hours} = 3648 \text{ seconds, until } 3648 / 204 = 17.9 \sim 18 \text{ patients month}$$

DISCUSSION


Pediatric dentistry's main objective in pulp treatment is maintaining the integrity and health of dental tissues. This is accomplished through the use of techniques and/or medications that allow the continuation of normal development until exfoliation and respect the particular characteristics of the life cycle of these elements.⁷

Comprehensive endodontic therapy for primary teeth can be challenging due to the peculiar anatomy of these teeth, as well as the child's behavior. For this reason, the length of the root must be correctly determined to minimize the apical periodontitis and possible damage to the permanent successor.³


Procedures applied in this study demonstrated that modern endodontic techniques had identical effects in reducing treatment time. In addition, this can limit the risk of clinical stress as much as possible.

The rotary technique required significantly less time compared to manual instruments.^{4,6,8} This is consistent with observations from other researchers who have noted that the optimization of endodontic treatment in primary teeth is important because it improves treatment quality and decreases clinical time.^{3,6,9-13} The principles of rotary instrumentation for deciduous teeth are the same as for permanent teeth.


The rotary instruments offer several advantages over traditional stainless steel instruments. They are flexible, have more cutting power, provide better maintenance of the original canal shape, considerably reduce deviation tendency or movement of the foramen, and reduce operative time.¹¹



Graph 1: Required time for instrumentation for the three systems



Graph 2: Percentage of time saved between systems (1 month)



Graph 3: Percentage of time saved between the two systems (12 months)

Several studies have been conducted comparing the Ni-Ti instruments' reciprocating and rotational movement. In evaluating the cyclic fatigue and bending of these instruments, there was a greater resistance observed for files applied in a reciprocal motion relative to the conventional rotational method, longer life-spans for these instruments, and a greater ability to maintain the centralized channel. In addition, the instrument's reciprocating movement caused less movement of apical foramen and less extrusion of dentin debris to the periapex.¹⁴⁻¹⁶ Results are consistent with the study showing that preparation time with rotary instruments was significantly less than with manual instrumentation, which is a clinically relevant factor for endodontic treatment.

CONCLUSION

Technological advances have simplified endodontic procedures with regard to the roundabout technique and reciprocation, even in the context of primary dentition, confirming other findings in the literature. However, good treatment also depends on the reduction or elimination of the infectious agent, appropriate instrumentation, efficient irrigation, and shutter-compatible antibacterial materials, as well as knowledge of the case study. However, well-designed follow-up studies for further statistical investigation are necessary for performing endodontic treatment of primary teeth.

REFERENCES

1. Braga MM, Oliveira LB, Bonini GA, et al. Feasibility of the International Caries Detection and Assessment System (ICDAS-II) in epidemiological surveys and comparability with standard World Health Organization criteria. *Caries Res* 2009;43(4):245-259.
2. Vasconcelos Cunha Bonini GA, Marcenos W, Oliveira LB, et al. Trends in the prevalence of traumatic dental injuries in Brazilian preschool children. *Dent Traumatol* 2009;25(6): 594-598.
3. Mello-Moura AC, Moura-Netto C, Araki AT, et al. Ex vivo performance of five methods for root canal length determination in primary anterior teeth. *IntEndod J* 2010;43:142-147.
4. Silva LA, Leonardo MR, Nelson-Filho P, Tanomaru JM. Comparison of rotary and manual instrumentation techniques on cleaning capacity and instrumentation time in deciduous molars. *J Dent Child* 2004;71:45-47.
5. Sasaki EW, Versiani MA, Perez DEC, et al. Ex vivo analysis of the debris remaining in flattened root canals of vital and non-vital teeth after biomechanical preparation with NiTi rotary instruments. *Braz Dent J, Ribeirão Preto* 2006;17(3):233-236.
6. Kummer TR, Calvo MC, Cordeiro MM, et al. Ex vivo study of manual and rotary instrumentation techniques in human primary teeth. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2008;105:84-92.
7. Corrêa FNP, Corrêa JP, Pellegrinetti MB, et al. Tratamento endodôntico em antecessor de dente de Turner. *Rev Inst Ciênc Saúde* 2008;26(2):258-262.
8. Nagaratina PJ, Shashikirann D, Subbareddy VV. In vitro comparison of Ni-Ti rotary instruments in root canal preparations of primary and permanent molar. *J Indian Soc Pedog Prev. Dent* 2006;24(4):186-191.
9. Barr ES, Kleier DJ, Barr NV. Use of nickel-titanium rotary files for root canal preparation in primary teeth. *Pediatr Dent* 1999; 21:453-454.
10. Walia HM, Brantley WA, Gerstein H. An initial investigation of the bending and torsional properties of Nitinol root canal files. *J Endod*, 1988;14:346-351.
11. Ferraz CC, Gomes NV, Gomes BP, et al. Apical extrusion of debris and irrigants using two hand and three engine-driven instrumentation techniques. *IntEndod J* 2001;34:354-358.
12. Pettiette MT, Delano EO, Trope M. Evaluation of success rate of endodontic hand files. *J Endod* 2001;27:124-127.
13. Moura ACVM, Borelli T, Matos R, et al. Como podemos otimizar a endodontia em dentes decíduos? Relato de caso. *Rev Assoc Paul Cir Dente* 2013;67(1):50-55.
14. Kuhn WG, Carnes DL Jr, Clement DJ, et al 3rd. Effect of tip design of nickel-titanium and stainless steel files on root canal preparation. *J Endod*. 1997;23:735-738.
15. Leonardo MR, Silva LAB, Nelson-Filho P, et al. Ex vivo evaluation of the accuracy of two electronic apex locators during root canal length determination in primary teeth. *Int Endodontic J*. 2008;41(4):317-321.
16. Kazemi RB, Stenman E, Spangberg LS. Machining efficiency and wear resistance of nickel-titanium endodontic files. *Oral Surg Oral Medi Oral Path Oral Radio and Endodontics* 1996;8:596-602.